

Tropical Cyclone Rainfall

Michael Brennan

National Hurricane Center



Outline


- Tropical Cyclone (TC) rainfall climatology
- Factors influencing TC rainfall
- TC rainfall forecasting tools
- TC rainfall forecasting process



Tropical Cyclone Rainfall Climatology



Tropical Cyclone Tracks









Global Mean Monthly TC Rainfall During the TC Season and Percent of Total Annual Rainfall

Data from TRMM 2A25 Precipitation Radar from 1998-2006








Contribution to Total Rainfall from TCs

- What percentage of average annual rainfall in southern Baja California, Mexico comes from tropical cyclones?
- 1. 10-20%
- 2. 20-30%
- 3. 40-50%
- 4. 50-60%

Khouakhi et al. (2017) *J. Climate*






Contribution to Global Rainfall from TCs (1970-2014 rain gauge study)


- Globally, highest TC rainfall totals are in eastern Asia, northeastern Australia, and the southeastern United States
- Percentage of annual rainfall contributed by TCs:
 - 35-50%: NW Australia, SE China, northern Philippines, Baja California
 - 40-50%: Western coast of Australia, south Indian Ocean islands, East Asia, Mexico

Khouakhi et al. (2017) *J. Climate*




Contribution to Global Rainfall from TCs (1970-2014 rain gauge study)





Contribution to Global Rainfall from TCs (1970-2014 rain gauge study)




- Relative contribution of TCs to extreme rainfall
- Gray circles indicate locations at which TCs have no contribution to extreme rainfall

Khouakhi et al. (2017) *J. Climate*



Annual TC Rainfall



- TC rainfall makes up a larger percentage of total rainfall during years when global rainfall is low
- Asymmetric generally more TC rainfall in the Northern Hemisphere
 - TCs produce 10-17% of global rain from 15-25°N
 - TCs produce 5-10% of global rain from 15-25°S



Frank Marks (HRD)

Biggest TC Rain Producers By Country/Island

Belize	829.8 mm	32.67"	Keith (2000)
Bermuda	186.7 mm	7.35"	October 1939 Hurricane
Canada	302.0 mm	11.89"	Harvey (1999)
Cayman Islands	764.8 mm	31.29"	Sanibel Island Hurricane (1944)
Costa Rica	920.0 mm	36.22"	Cesar (1996)
Dominica	422.3 mm	16.63"	Jeanne (2004)
Dominican Rep.	1001.5 mm	39.43"	Flora (1963)
El Salvador	406.4 mm	16.00"	Adrian (2005)
Guadeloupe	508.0 mm	20.00"	Marilyn (1995)
Guatemala	600.0 mm	23.62"	Mitch (1998)
Haiti	1447.8 mm	57.00"	Flora (1963)
Honduras	912.0 mm	35.89"	Mitch (1998)
Martinique	680.7 mm	26.80"	Dorothy (1970)
Mexico	1576.0 mm	62.05"	Wilma (2005)
Nicaragua	1597.0 mm	62.87"	Mitch (1998)
Panama	695.0 mm	27.36"	Mitch (1998)
Puerto Rico	1058.7 mm	41.68"	T.D. #19 (1970)
St. Lucia	668.0 mm	26.30"	Tomas (2010)
St. Martin/Maarten	866.6 mm	34.12"	Lenny (1999)
Venezuela	339.0 mm	13.30"	Brett (1993)




Original Source: David Roth WPC (2006)

Biggest TC Rain Producers By Country/Island

Belize	829.8 mm	32.67"	Keith (2000)	
Bermuda	186.7 mm	7.35"	October 1939 Hurricane	
Canada	302.0 mm	11.89"	Harvey (1999)	
Cayman Islands	764.8 mm	31.29"	Sanibel Island Hurricane (1	944)
Costa Rica	920.0 mm	36.22"	Cesar (1996)	
Cuba	2550.0 mm	100.39"	Flora (1963)	
Dominica	422.3 mm	16.63"	Jeanne (2004)	
Dominican Rep.	1001.5 mm	39.43"	Flora (1963)	
El Salvador	406.4 mm	16.00"	Adrian (2005)	
Guadeloupe	508.0 mm	20.00"	Marilyn (1995)	
Guatemala	600.0 mm	23.62"	Mitch (1998)	
Haiti	1447.8 mm	57.00"	Flora (1963)	
Honduras	912.0 mm	35.89"	Mitch (1998)	
Jamaica	3429.0 mm	135.00"	November 1909 Hurricane	
Martinique	680.7 mm	26.80"	Dorothy (1970)	
Mexico	1576.0 mm	62.05"	Wilma (2005)	
Nicaragua	1597.0 mm	62.87"	Mitch (1998)	
Panama	695.0 mm	27.36"	Mitch (1998)	
Puerto Rico	1058.7 mm	41.68"	T.D. #19 (1970)	
St. Lucia	668.0 mm	26.30"	Tomas (2010)	
St. Martin/Maarten	866.6 mm	34.12"	Lenny (1999)	
Venezuela	339.0 mm	13.30"	Brett (1993)	Hu



Original Source: David Roth WPC (2006)




Characteristics of TC Precipitation

Stratiform and Convective Mechanisms
Stratiform Rain ~50% of Total Rain from TC

NOAA/HRD - Daily Radar Rainfall Estimate Study


Typical warm season 1-day total


Hurricane Irene (1999) 1-day total



6

TC Rainband Complexes: Stratiform Region







- Broad stratiform band in left-of-shear half of the storm
- Mesoscale ascending outflow & descending inflow driven by latent heating & latent cooling patterns
- Increased rainfall along line where descending inflow halts
- Descending inflow strengthens the outer core of vortex



TC Rainband Complexes: Convective Cells





Cell 1

- Weaker, shallower reflectivity core
- Weaker updrafts
- Shallower, but stronger inflow layer
- Tangential jet and outflow confined to lower levels
- More intense reflectivity, heavier rain

Cell 2

- Increased CAPE, more buoyant updrafts
- Deeper inflow layer
- Tangential jet and outflow extend deeper into the troposphere



Factors Influencing Tropical Cyclone Rainfall





- Which of the following is NOT a primary factor in determining rainfall from tropical cyclones?
 - 1. TC Track
 - 2. TC Size
 - 3. TC Intensity
 - 4. Topography





What Factors Influence Rainfall from Tropical Cyclones?

- Movement slow forward motion can produce more rain
- Storm size the larger the storm, the greater the area typically receiving rain
- Storm track determines the location of the rain
- Diurnal cycle heaviest rainfall generally near the storm center overnight, outer band rainfall during the day
- Topography enhances rainfall in upslope areas, but decreases rainfall past the spine of the mountains
- Moisture entrainment of dry air can redistribute and/or reduce the amount of precipitation; increased moisture can increase rainfall
- Interaction with other meteorological features (troughs, fronts, jets) and extratropical transition can greatly modify rainfall distribution




Storm Motion

Slow vs. fast moving TCs

 TCs with a turning or looping track vs.
 straight mover Hurricane Mitch fatalities: Honduras: 5,677

Nicaragua: 2,863 Guatemala: 258

El Salvador: 239


































Factors Influencing TC Rainfall Storm Size

Determined by distance from center to outermost closed isobar

<2 degrees	"Very small/ midget"	Marco (2008)
2-3 degrees	"Small"	lda (2009)
3-6 degrees	"Average"	Frances (2008)
6-8 degrees	"Large"	Wilma (2008)
>8 degrees	"Very large"	Sandy (2012)



Factors Influencing TC Rainfall

Time of Day Alberto, July 4-5, 1994

















Factors Influencing TC Rainfall


Time of Day Alberto, July 4-5, 1994







00z












Terrain Impacts


Heaviest rainfall favors mountains perpendicular to the wind




28,005 homes destroyed

Factors Influencing TC Rainfall

Vertical Wind Shear – Northern Hemisphere




Shear directed across
the storm track leads
to more uniform
distribution of the
rainfall



Shear directed parallel
to the storm track
leads to a distribution
of the rainfall
asymmetry on the left
side of the track






Factors Influencing TC Rainfall

Environmental Steering in Northern Hemisphere

- Very slow moving TCs and symmetrical TCs produce the most rainfall near the center
 - Maximum rainfall at night (especially when over land)
 - Weak steering flow
- TCs that move into a break in the subtropical ridge often produce most of the rain *right* of their track




Even rainfall distribution = Rain over smaller area on right side = Higher totals

- TCs that recurve due to significant upper troughs in the westerlies often produce most of their rain *left* of their track
 - Rainfall may spread well in advance of the TC due to interaction with the upper jet on the leading edge of the trough


Factors Influencing TC Rainfall Predecessor Rainfall Events





- Moisture transport well ahead of TC itself
- Coherent area of rain displaced north of the TC (near a front or over terrain)
- Maximum rainfall rates can exceed 200 mm in 24 hr
- Occurs for approximately 1 of 3 landfalling TCs in U.S.





Where is Flooding from Tropical **Cyclones Likely to Occur?** Areas where the ground is already saturated (low flash flood guidance values) Valleys/watersheds Areas of orographic enhancement

- Areas with poor drainage or prone to runoff
- Areas with directed drainage that can be overwhelmed







TC Rainfall Forecasting Tools





NHC Rainfall Product


Incorporates microwave (MW) satellite data rainfall rates

- NHC uses two different merged satellite rainfall estimation techniques to replace the IR-only method:
 - NRL-Blend and QMORPH incorporate available MW data and propagate precipitation forward in time via IR
 - Training on the NRL-Blend technique: http://www.nrlmry.navy.mil/training-bin/training.cgi
- As a third product, NHC uses the last applicable GFS forecast
 - A model forecast has the advantage of dynamics, topography, moisture, etc.



NHC Rainfall Product: Why Microwave?


- Geostationary IR data provides excellent spatiotemporal resolution, but is not optimal for rain estimation
- Microwave provides improved rainfall accuracy but at low temporal resolution
- Quantitative precipitation estimate (QPE) products leverage each method's strength...




Satellite Rainfall Estimates

International Precipitation Working Group (IPWG) has an exhaustive list of data sources for precipitation information, some of it in real time.

http://www.isac.cnr.it/~ipwg/data/datasets.html



Satellite Rainfall Estimates International Precipitation Working Group (IPWG) has an exhaustive list of data sources for precipitation information, some of it in real time. http://www.isac.cnr.it/~ipwg/data/datasets.html

























NHC Rainfall Product: Text

TCCA22 KNHC 291843 STDWCA

SATELLITE TROPICAL DISTURBANCE RAINFALL ESTIMATES NUS NATIONAL HURRICANE CENTER MIAMI FL 2115 UTC TUE AUG 29 2009

SYSTEM NAME	DATE/TIME	LOCATION
I.S ANDRES	29/1800 UTC	17.2N 102.3W

RAINFALL ESTIMATED BY SATELLITE VIA OMORPH. . 24-HOUR RAINFALL MAXIMUM (FROM 18-18 UTC) - 235 MM AT 23.3N 99.2W 6-HOUR RAINFALL MAXIMUM (FROM 12-18 UTC)- 150 MM AT 24.2N 100.5W RAINFALL DISTRIBUTION IN MM OVER THE LAST 6-HOURS (FROM 12-18 UTC

LATITUDELONGITUDE									
		104W-	103W	103W-102W	102W-101W	101W-100W	100W- 99W	99W- 98W	
	27-28N	0-	40	5- 40	10- 45	10- 45	10- 30	0- 20	
	26-27N	5-	40	10- 45	15- 55	20- 50	15- 30	5- 20	
	25-26N	15-	45	20- 70	35- 85	60-100	30- 70	20- 45	
	24-25N	40-	76	55-100	100-130	110-150	60-100	40- 75	
	23-24N	20-	50	45- 70	70- 90	70- 95	40- 65	15- 40	
	22-23N	0-	35	5- 40	10- 30	10- 25	5- 25	0- 10	

24-HOUR RAINFALL MAXIMUM (FROM 18-18 UTC) - 295 MM AT 23.3N 98.7W 6-HOUR RAINFALL MAXIMUM (FROM 12-18 UTC)- 125 MM AT 24.6N 100.2W RAINFALL DISTRIBUTION IN MM OVER THE LAST 6-HOURS (FROM 12-18 UTC).

ı	LATITUL						
ı		104W-103	W 103W-102W	102W-101W	101W-100W	100W- 99W	99W- 98W
ı	27-28N	0- 35	5- 40	10- 45	10- 45	5- 25	0- 20
ı	26-27N	0- 35	10- 45	15- 50	20- 50	10- 30	5- 20
ı	25-26N	15- 45	20- 70	35- 80	65-100	25- 70	15- 45
ı	24-25N	35- 75	55- 95	100-120	110-125	60-100	35- 75
ı	23-24N	20- 45	45- 75	65- 85	70- 95	35- 70	15- 40
ı	22-23 N	0- 30	5- 40	10- 30	10- 30	5- 25	0- 10
ı							

RAINFALL HINDCAST FROM THE 06Z GFS MODEL.

24-HOUR RAINFALL MAXIMUM (FROM 18-18 UTC)- 305 MM AT 23.1N 101.8W 6-HOUR RAINFALL MAXIMUM (FROM 12-18 UTC)- 130 MM AT 24.9N 101.9W RAINFALL DISTRIBUTION OVER THE LAST 6-HOURS (FROM 12-18 UTC)...

LATITU	DE		LONGITUDE.				
	.104W-	103W	103W-102W	102W-101W	101W-100W	100W- 99W	99W- 98W
27-28N	r o-	30	5- 40	10- 45	15- 45	5- 25	0- 20
26-27N	-0	35	10- 45	15- 45	20- 50	10- 30	5- 20
25-26N	15-	45	20- 70	35- 85	60-100	30- 70	20- 45
24-25N	35-	75	55-100	100-130	100-125	65-100	40- 75
23-24N	20-	45	45- 70	70- 85	70- 95	40- 70	15- 40
22-23N	T 5-	35	5- 40	10- 30	10- 25	5- 25	0- 10

ESTIMATES INDICATE UNCERTAINTY IN THE AMOUNT OF PAIN RECEIVED

RAINFALL MAY BE UNDERESTIMATED ON THE WINDWARD SIDE OF TERRAIN

FOR ADDITIONAL INFORMATION PLEASE VISIT HTTP://WWW.HURRICANES.GOV/RAINFALL

FORECASTER NELSON

Created for every "invest" system

Can be created for any disturbance

Rainfall product still available in text format like the old product

Differences in content and format compared to the old product:

- 6-hour quantitative precipitation estimates from 3 methods
 - Presented as a range of rainfall within a 10x10 box
 - Covers total area of 6°x6° centered near disturbance
- Earth-relative coordinates (i.e. no reference to "left-of-center"/"right of center")
- Available at: http://www.nhc.noaa.gov/marine/rainfall/



New NHC Rainfall Product: Text

TCCA22 KNHC 291843 STDWCA

NUS NALLONAL HUM TOLL 2115 UTC TUE AUG 29

SYSTEM NAME T.S ANDRES

RAINFALL ESTIMATED B' 24-HOUR PAIN ALL MAXI 6-HOUR RAIN ALL MAXIN RAINFALL DISTRIBUTION

LATITUDF/104W-103W 103T 0- 40 27-28N 26-27N 5- 40 15- 45 24-25N 40- 76 2**%**-24N 20- 50 22-23N 0 - 35

RAINFALL ESTIMATED B' 24-HOUR PAINFALL MAX 6-HOUR RAINFALL MAXIN RAINFALL DISTRIBUTION

LATITUDE.....104W-103W 103U

22-23N

27-28N 0 - 350- 35 26-27N 25-26N 15- 45 24-25N 35- 75 55- 95 100-120 23-24N 20- 45 45- 75

0 - 30

RAINFALL ESTIMATED BY SATELLITE VIA NRL-BLEND...

24-HOUR RAINFALL MAXIMUM (FROM 18-18 UTC)- 295 MM AT 23.3N 6-HOUR RAINFALL MAXIMUM (FROM 12-18 UTC)- 125 MM AT 24.6N 100.2W RAINFALL DISTRIBUTION IN MM OVER THE LAST 6-HOURS (FROM 12-18 UTC)...

LATITUDE

	1040-1030	1030-1020	1020-1010	TOTM-TOOM	TOOM- 33M	99W- 98W
27-28N	0- 35	5- 40	10- 45	10- 45	5- 25	0- 20
26-27 N	0- 35	10- 45	15- 50	20- 50	10- 30	5- 20
25-26 N	15- 45	20- 70	35- 80	65-100	25- 70	15- 45
24-25N	35- 75	55- 95	100-120	110-125	60-100	35- 75
23-24 N	20- 45	45- 75	65- 85	70- 95	35- 70	15- 40
22-23 N	0- 30	5- 40	10- 30	10- 30	5- 25	0- 10

RAINFALL HINDCAST FROM THE 06Z GFS MODEL...

24-HOUR RAINFALL MAXIMUM (FROM 18-18 UTC) - 305 MM AT 23.1N 101.8W 6-HOUR RAINFALL MAXIMUM (FROM 12-18 UTC)- 130 MM AT 24.9N 101.9W RAINFALL DISTRIBUTION OVER THE LAST 6-HOURS (FROM 12-18 UTC)...

65-85

10- 30

110-125

10- 30

70- 95

35- 70

5- 25

15- 40 0 - 10

	.104W-103W	103W-102W	102W-101W	101W-100W	100W- 99W	99W- 98W
27-28N	0- 30	5- 40	10- 45	15- 45	5- 25	0- 20
26-27N	0- 35	10- 45	15- 45	20- 50	10- 30	5- 20
25-26N	15- 45	20- 70	35- 85	60-100	30- 70	20- 45
24-25N	35- 75	55-100	100-130	100-125	65-100	40- 75
23-24N	20- 45	45- 70	70- 85	70- 95	40- 70	15- 40
22-23N	5- 35	5- 40	10- 30	10- 25	5- 25	0- 10

DIFFERENCES BETWEEN THE SATELLITE AND MODEL DERRIVED PAINFALL ESTIMATES INDICATE UNCERTAINTY IN THE AMOUNT OF RAIN RECEIVED

RAINFALL MAY BE UNDERESTIMATED ON THE WINDWARD SIDE OF TERRAIN

FOR ADDITIONAL INFORMATION PLEASE VISIT HTTP://WWW.HURRICANES.GOV/RAINFALL

FORECASTER NELSON

- Lat-lon grid of rainfall accumulation
- 6-h accumulation ranges (in mm)
- Differences in the 3 rainfall estimates reveal uncertainty
- Available at: http://www.nhc.noaa.gov/marine/rainfa




Ensemble Tropical Rainfall Potential Product (eTRaP)

6-hourly Day 1 forecasts: Extrapolates polar orbiting satellite rain rate along TC forecast tracks

(AMSU, SSMI, AMSR, GPM)

- A satellite "member" is included when its path passes over the
- "Members" are weighted according to age of pass and past performance of sensor
- Official forecast of TC track & at least 2 members needed to create a forecast
- Updated daily at 0315, 0915, 1515, and 2115 UTC



Forecast Period

Probability exceedance


Quantitative Precipitation Forecast

http://www.ssd.noaa.gov/PS/TROP/etrap.html





eTRaP: http://www.ssd.noaa.gov/PS/TROP/etrap.html Probability of **Probability of Probability of** > 25 mm > 100 mm > 50 mm Typhoon Ului 06-12 hr eTRaP forecast Ø



CLIQR: Picking an Analog for a TC Rainfall Event

www.wpc.ncep.noaa.gov/tropical/rain/web/cliqr.html

Looks at:

- The current rain shield size and compare it to TCs from the past
- How fast is the TC moving?
- Vertical wind shear in current/past events?
- Look for storms with similar or parallel tracks
- Is topography a consideration?
- Look for nearby fronts and examines the depth of nearby upper troughs for current event and possible analogs

Not all TC events will have a useful analog





Tropical Cyclone Rainfall Data

http://www.wpc.ncep.noaa.gov /tropical/rain/tcrainfall.html

Tropical Cyclone Rainfall Data

Rainfall analogs to current tropical

cyclones

Select Storm By Point Of Entry

Point Maxima for Tropical Cyclones

Tropical Cyclone Rainfall Slideshow (in

Powerpoint format)



Select Storm By Name

Select Storm By Region Of Impac

Tropical Cyclone Maxima Per

Mexican State

Tropical Cyclone Rainfall

Forecasting Acknowledgments This page is under construction...so new information will be added as time allows. Data is available for tropical and subtropical cyclones that impacted the U.S. from 1963 onward to the present, and Mexico between 1995 and 2003, as well as some older historic storms. The image of Hurricane Floyd shown to the left was provided by the Operational Satellite Events Imagery web page of NOAA. Please select the page of your choice from the following list.

CLIQR Matching TC List (Rainfall Matches Accessible via Hyperlink)

For any questions, comments, suggestions, e-mail David.Roth@noaa.gov Last updated May 26, 2009



Results ranked from best match to worst match, with ties being won by the earlier storm. BETA 2005: No graphic available.

HATTIE 1961: No graphic available.

JOAN 1988

MARCO 1996: No graphic available. NOT NAMED 1964: No graphic available.


GORDON 1994 KATRINA 1999

MARTHA 1969: No graphic available. THIRTEEN 1985: No graphic available. BRET 1993: No graphic available.

ALMA 1970

IRENE 1971: No graphic available. UNNAMED 1981: No graphic available. FOURTEEN 2002: No graphic available. SIX 1969: No graphic available. LAURA 1971: No graphic available. SEVENTEEN 1973: No graphic available. CESAR 1996: No graphic available.

Available for active TCs at: www.wpc.ncep.noaa.gov/tropical/ rain/web/cliqr.html






Select Storm By Year

Tropical Cyclone Maxima Per U. S.

Tropical Cyclone Averages and

Maxima per Duration

Methodology for climatology



Launched in October 2016

Rainfall Rate Algorithm

Rainfall Potential Algorithm


Probability of Rainfall Algorithm








Rainfall Rate



- The algorithm generates
 estimates of the instantaneous
 rainfall rate at each IR pixel
- Uses IR brightness temperatures and is calibrated in real time against microwavederived rain rates to enhance accuracy
- The higher spatial and temporal resolution available from GOES-R will be able to automatically resolve rainfall rates on a finer scale


Rainfall Potential



- Predicted rainfall accumulation for the next 3 hours at the satellite pixel scale
- Extrapolation from current and previous rainfall rates from the GOES-R Rainfall Rate Algorithm




Probability of Rainfall




- Generates a gridded probability of at least 1 mm of rainfall during the next 3 hours at the satellite pixel scale
- Uses intermediate rainfall rate forecasts from the Rainfall Potential Algorithm as input to a statistical model calibrated against estimates from the Rainfall Rate Algorithm














NCEP Model QPF Biases

- NCEP models are updated frequently which makes it difficult to isolate distinct biases
- Run-to-run consistency increases confidence of occurrence
- Convective feedback problems
 - Updraft overtakes the grid cell
 - WPC estimates that QPF maximum amounts are reasonable about half the time when convective feedback is noted, but the location can be far off (Roth, 2008)






Model TC QPF Skill Marchok et al, 2007



1998-2004 U.S. landfalling TC QPFs from the GFS, GFDL hurricane model, the NAM, and the R-CLIPER (Rainfall Climatology and Persistence)


Three elements of TC rainfall forecasts used as a basis for comparing models:

- model ability to match the large-scale rainfall pattern,
- model ability to match the mean rainfall and the distribution of rain volume, and
- model ability to produce the extreme amounts often observed in TCs



Model TC QPF Skill Marchok et al, 2007

- Compared to R-CLIPER, all numerical models showed comparable or greater skill for all attributes
- GFS performed the best of all of the models for each of the categories
- GFDL had a bias of predicting too much heavy rain, especially in the core of the tropical cyclones
- NAM predicted too little of the heavy rain.
- R-CLIPER performed well near the track of the core, but predicted much too little rain at large distances from the track





Model TC QPF Skill Marchok et al, 2007

- Compared skill for all a
- GFS perfor
- GFDL had a the tropical
- NAM predic
- R-CLIPER little rain at


Track forecast error was a primary determinant of tropical cyclone **QPF** error

e or greater


ories

he core of

much too







Where to Find Model QPFs

NHC Tropical Rainfall Webpage (storm-specific GFS, hindcasts)

http://www.nhc.noaa.gov/marine/rainfall/

 NCEP models (GFS, NAM, GEFS, NAEFS) including tropical guidance (HWRF and HMON)

http://mag.ncep.noaa.gov

Canadian Global GEM

http://www.weatheroffice.gc.ca/model_forecast/global_e.html

Canadian Global GEM Ensembles

http://www.weatheroffice.gc.ca/ensemble/index_e.html

NAVGEM

http://www.nrlmry.navy.mil/metoc/nogaps/

ECMWF

http://schumacher.atmos.colostate.edu/weather/ecmwf.php

Penn State Tropical Atlantic E-Wall

http://mp1.met.psu.edu/~fxg1/ewalltropatl.html




TC QPF Forecast Process



NWS Tropical Cyclone Quantitative Precipitation Forecasts (QPF) Michita Falls 1.33 • ♥ 0.70 Fort Horth Dallas Terrell o 0.46 o () d) Palestine · 0.07 WFO QPF 9.00 -8.00 -7.00 -6.00 -5.00 -4.00 -3.00 -2.50 -2.00 -1.75 ecip.Amt(in) Ending Wed Apr 29 2009 7PM CDT (Thu Apr 30 2009 00Z) NWS Fort Worth, TX Graphic Created Apr 29 2:49PM CDT WPC QPF

Production of Tropical Cyclone Quantitative Precipitation Forecasts A good place to start is the model closest to the NHC track forecast Day 1 QPF 24-h ending 12Z 21 Aug 2008 -T.S. Fay 9.00






9.00 8.00 1.00

MELBOURNE FLORIDA.


Production of Tropical Cyclone Quantitative Precipitation Forecasts

How well do the models match the NHC rainfall statement?



Production of Tropical Cyclone Quantitative Precipitation Forecasts

Use observations and recent model data to determine the current structure/rainfall rates






Servicio Hidrologico Nacional

Production of Tropical Cyclone Quantitative Precipitation Forecasts

Locate relevant synoptic scale and meso-scale boundaries




Tropical Surface Analysis Mon 18:00Z 24-Oct-11

MIMIC Total Precipitable Water (mm)



Production of Tropical CycloneQuantitative Precipitation Forecasts

Use conceptual models and pattern recognition as well as the forecast upper level winds to further adjust QPF





Identify areas of orographic enhancement



























Identify areas of orographic enhancement





Production of Tropical Cyclone Related QPF

Determine how a change in available moisture could increase, decrease, or redistribute rainfall




























Production of Tropical Cyclone Related QPF

Use climatology (CLIQR, R-CLIPER, TC Rainfall Climatology) and data from past storms to:

- Increase/decrease amounts
- Adjust numerical guidance biases
- Reality check
- Highlight areas significantly impacted by terrain effects

INVEST_AL96

Results ranked from best match to worst match, with ties being won by the earlier storm. BETA 2005: No graphic available.

GERT 1993

HATTIE 1961: No graphic available.

JOAN 1988


MARCO 1996: No graphic available. NOT NAMED 1964: No graphic available.

GORDON 1994 KATRINA 1999

MARTHA 1969: No graphic available. THIRTEEN 1985: No graphic available. BRET 1993: No graphic available.

<u>ALMA 1970</u>

IRENE 1971: No graphic available.
UNNAMED 1981: No graphic available.
FOURTEEN 2002: No graphic available.
SIX 1969: No graphic available.
LAURA 1971: No graphic available.
SEVENTEEN 1973: No graphic available.
CESAR 1996: No graphic available.








In Conclusion

- Remember the factors that influence TC rainfall (size of storm, time of day, speed etc.)
- Evaluate the quality of the model data available to you compared to the current conditions
- Assess the amount of shear in the environment How will it influence rainfall?
- Are there past TCs that resemble the rainfall distribution and forecast of the TC?
- Use all of the tools available (satellite rainfall products, NWP models, etc.)
- Remember, heavy rain can also occur well away from the TC itself (PRE, secondary disturbances, etc.)















Thank You

Questions?

