



Ensemble Prediction Systems



Eric Blake

National Hurricane Center

March 1, 2023

Acknowledgements to Dr. Michael Brennan

Question 1

What are some current advantages of using single-model ensembles?


- A. Estimates of uncertainty
- B. TC intensity model spread
- C. Alternative TC-track solutions
- D. All of the above
- E. A & C

Why Aren't Models Perfect?

- Atmospheric variables cannot be measured to an infinite degree of accuracy or precision (measurement error)
- Models' initial state never matches the real atmosphere (analysis error)
- Initial condition errors grow with model integration time, most rapidly at smaller scales (error growth)
- Model equations do not fully represent all of the processes in the atmosphere (model error)
- Model grid cannot explicitly resolve all features and processes in the atmosphere (model error)

How did we manage to extend the FSH beyond 2 weeks?

Predictable signals propagate from the better-initialized and more predictable scales ('mainly' the large scales, the slowly evolving components) to the less predictable (small/fast) scales



Errors propagate from poorly initialized scales ('mainly' the smaller scales) thus reducing the predictive skill

(Buizza and Leutbecher 2015, QJRMS)

Options?

- Increase our understanding of physical processes and how models represent them (research)
- More accurate and numerous observations with greater coverage (expensive)
- Improved data assimilation methods (4-D Variational Data Assimilation, Ensemble Kalman Filter)
- Faster computers and more complex models (many programs competing for resources)
- *Probabilistic forecasting with ensembles*

Definitions


- **Deterministic Model** - single forecast from one forecast model or method using a single set of initial conditions
 - Examples: GFS, ECMWF, UKMET, HMON, HWRF, HAFS
- **Ensemble** - collection of “member” forecasts verifying at the same time created from:
 - Different but equally viable initial conditions
 - Different forecasting methods and/or models that (ideally) statistically represent nearly all forecast possibilities

Definitions

- **Dynamical Model Ensemble** –based on perturbation of initial conditions of a single model or different models to create “member” forecasts
 - Examples: NCEP Global Ensemble Forecast System (GEFS), ECMWF Ensemble Prediction System (EPS)
- **Control Run** – for dynamical model ensembles, usually the member of the ensemble run with the “best” initial analysis
 - The analysis used by the control run is usually perturbed to produce initial conditions for the remaining ensemble members
- **Spread** – measure of the degree of disagreement (i.e., standard deviation) between ensemble members



Ensemble Use

- Originally used for medium-to long-range forecasting of the large-scale pattern
- Uses have grown to encompass all temporal and spatial scales down to convective storm scale
- Address uncertainty, particularly those leading to rapidly diverging solutions
 - Initial conditions, model physics, resolution, model numerics



Ensemble Use

- Estimate rate of skill loss with time
 - Spread of solutions generally increases with time
- Compute probabilities for a particular event or condition
 - 25 mm of precipitation, winds > 34 kt
- Identify regions where the analysis and forecast are sensitive to additional data in the analysis
 - Ensemble Kalman Filter, targeted observations (next week Ryan Torn's talk)




Ensemble Mean vs. Deterministic

- Deterministic runs (e.g., GFS) usually have more skill than any *individual ensemble member* due to superior resolution
- Ensemble mean usually has at least as much skill as an *equal-resolution control run*
- Ensemble mean can be more skillful than a *higher-resolution deterministic run*, especially beyond ~3 days

Current Global Ensemble Systems that NHC uses most frequently

NCEP Global Ensemble Forecast System (GEFS)

- 4 cycles per day (00, 06, 12, 18 UTC)
- 31 members (1 control + 30 perturbed)
- Forecast extends out to 384 hours (16 days)




180-h forecast of 588 dm 500-mb height contour valid at
1200 UTC 22 March 2010


NCEP GEFS

- Current Configuration (last upgrade Sep 2020)
 - ~25 km resolution, run to 35 days at 00Z
 - 64 vertical levels
- Ensemble members
 - 30 members generated using EnKF
 - Uses stochastically perturbed physics tendencies (SPPT) scheme and stochastic kinetic energy backscatter (SKEB) scheme for perturbations
 - Model physics consistent with GFS
- Deterministic GFS (2024 upgrade coming?)
 - ~13 km resolution for full run (16 days)
 - 127 vertical levels

GEFS Mean vs. GFS (2019-2022)



In the Atlantic, the GEFS ensemble mean track forecast (**AEMI**) is competitive with the deterministic GFS (**GFSI**) through 36 h and better afterward



Eastern Pacific – ensemble better at all times

ECMWF Ensemble Prediction System

- 51 members (1 control+50 perturbed members)
- Run twice daily (00 and 12 UTC) out to 15 days, 6/18 UTC 144h
 - T639 (~ 18 km) to 15 days
 - 137 vertical levels

Perturbations:

- Generated using singular vectors and stochastically Perturbed Parameterization Tendencies Scheme (SPPT)

Deterministic ECMWF

- Horizontal grid resolution T1279 (~9 km) out to 10 days with 137 vertical levels

Big ensemble upgrade coming in June for horizontal resolution to match deterministic!



“Good” and “Bad” Ensembles

An ensemble forecast starts from initial perturbations to the analysis...

In a good ensemble “truth” looks like an member of the ensemble
(Toth, 1992)


The initial perturbations should reflect the analysis “errors of the day”

A bad ensemble is still useful (implies there is a bug in the system)



Kalnay 2019

Matthew ensemble guidance 1 Oct 00 UTC




Old GEFS pre-2020 (blue) too underdispersive, especially in Caribbean

Every single GEFS member also too fast at 5 days

ECMWF (red) has more realistic spreads, albeit potentially too large

ECMWF ensemble tracks for Laura at Aug 25 0000 UTC




Every member was too far west

Don't just worship at the altar of the ECMWF!


Ensemble Display and Interpretation

Displaying Ensembles



If we try to look at every ensemble member at once, it is messy and difficult to interpret


Displaying Ensembles



Ensemble Mean - average of multiple forecast members verifying at same time

Displaying Ensembles

Ensemble Mean and Spread



- **Black** lines = ensemble mean 500-mb height forecast
- Spread indicated by shading (meters)
 - **Orange/Red** – little agreement between members
 - **Blue** – good agreement between members

Displaying Ensembles


Ensemble Mean and Spread

- Advantages
 - Summarizes data in easy to interpret form
 - Information provided for the entire domain
 - Low predictability features smoothed out by the ensemble mean and easily identifiable using spread
- Disadvantages
 - Ensemble mean can be misleading (and may not be the best forecast) if multiple clusters of nearly equal probability forecast outcomes exist (i.e., bi-modal distribution)
 - May not reveal extreme outlier solutions

Interpreting Mean and Spread

Large spread within the ensemble mean feature ? **Uncertainty in amplitude of the feature**


- In this case, there is uncertainty in the **depth** (not the location) of this 500-mb trough
- If there were a tropical cyclone located southeast of this trough, would the trough be deep enough to recurve the tropical cyclone?



Interpreting Mean and Spread

Large spread upstream or downstream of an ensemble mean feature ? **Uncertainty in the location of the feature**


- In this case, there are nearly equal chances that the 500-mb trough will be east or west of the position shown by the ensemble mean trough
- If a tropical cyclone was located southeast of this trough, at what time will the tropical cyclone begin to be influenced by this trough?



Interpreting Mean and Spread

Large spread on one side of an ensemble mean feature ? **A cluster of ensemble members different from the ensemble mean**

- In this case, the spread indicates greater potential for the trough axis to be east of the ensemble mean trough than to the west
- If there was a tropical cyclone located southeast of this trough, at what time will the tropical cyclone begin to be influenced by this trough?



Single-Model Ensembles for TC Track Forecasting


NCEP Global Ensemble Forecast System

Tropical Cyclone Track Forecast Guidance




ECMWF Ensemble


Sandy example of desirable spread/verification




Joaquin ensemble guidance




GFS Joaquin ensembles 29 Sep 1200 UTC



ECMWF Joaquin ensembles 29 Sep 1200 UTC




Track dependent on intensity?



-Stronger members farther right
-Weaker members farther north

Ensemble Problems


- Need a properly calibrated system
 - GEFS used to be underdispersive (much better since upgrade)
 - This problem results in an overconfident forecast (UKMet ensemble has this more than EPS/GEFS now)
 - Lower resolution can also hinder a more accurate track forecast (i.e. when track especially dependent on intensity)
- Other issues
 - Ensemble mean can be misleading (and may not be the best forecast) if multiple clusters of nearly equal probability forecast outcomes exist (i.e., bi-modal distribution)
 - May not reveal extreme outlier solutions




Hurricane Leslie

Long lasting and *highly* annoying

Leslie track guidance 10/9/6z



Major model spread, but best models farthest south




Leslie track guidance 10/10/6z



EC
ensemble
not as
definitive

ECMWF Ens. (0-120h only), init: 2018101000, AL13 Leslie color = max wind (kt)




Leslie track guidance 10/10/12z



72 hour zoom


—
All models have a
400- 600 n mi error

Leslie track guidance 10/10/18z




Few more models
shift north, but
most still miss the
trough

EC ensemble
continues to
shift south




Leslie track guidance 10/12/6z




Models mostly
jump northward


ECMWF Ens. (0-120h only), init: 2018101200, AL13 Leslie color = max wind (kt)



Leslie track guidance 10/12/18z




Finally!




Track Forecast Challenges - Marco

Potential interaction between Marco and Laura as well as uncertainty about Marco's intensity led to huge variability in track guidance for Marco and poor forecasts



Marco track guidance 21-23 August 2020


Incorrect Model Trends Near Landfall - Laura



Laura guidance 1800 UTC Aug 24

Guidance almost perfectly centered around the Louisiana landfall

Incorrect Model Trends Near Landfall - Laura




Laura guidance 0000 UTC Aug 25

6 hours later- trouble.

Notable westward shifts of
ECMWF ensemble and
corrected-consensus aids

Incorrect Model Trends Near Landfall - Laura



Laura guidance 0600 UTC Aug 25


Lots of models shifted west

NHC fcst did not leave LA

Expectation of a deep hurricane/SW winds in high-levels led NHC to stay on the E side of the guidance

Huge Houston evacuation implications

Track/Warning Forecast Challenges - Sally




Track guidance 0600 UTC 13 September 2020

Track guidance whiffed when the Hurricane Warning was put up

Resulted in a clear-sky bust for Louisiana

Note HMNI closest - yet it had a NE bias for Marco. Recent past does affect forecaster perceptions

Track/Warning Forecast Challenges - Sally



ECMWF Ens 0600 UTC 13 September 2020


No help from the best ensemble system either

If anything you might suspect the forecast would bust left, not right

Some stronger members on right side, but intensity skill lags

Ian Super-ensemble (EPS + GEFS + UKMET + CMC)
Track Density (350-km radius)

Hour 96 | Valid 0000 UTC 29 September 2022
Initialized 0000 UTC 25 September 2022



Wave of the future – put all systems together and use probabilities?

Genesis Guidance

Little objective guidance is seen with ensembles now, though they help subjectively.

In-house product?

shading: combined probability of 70 ensemble members (GEFS + ECENS):

- 850 – 700 hPa RH > 70%
- 200 – 850 hPa vertical wind shear < 20 kt

contours: 850 hPa relative vorticity ($8 \times 10^{-5} \text{ s}^{-1}$ intervals)

thin green: ECENS members


thick green: ECMWF deterministic

thin yellow: GEFS members

thick yellow: GFS deterministic

Invest AL93


0000 UTC November 6, 2015 + 66 h



TC Intensity Ensemble Forecasting

- Some skill above single-model deterministic
- Very computational expensive to run high-resolution (<3 km) intensity ensembles
- HFIP is funding efforts to find products that could be operationally useful

TC = 09L, DTG = 2021082612



ECWMF experiment, 2030 desired resolution


IRMA operational v. 5km

Richardson,
ECMWF, 2018


Date 20170905 12 UTC @ ECMF

Probability that **IRMA** will pass within 120 km radius during the next 240 hours
tracks: **solid**=HRES; **dot**=Ens Mean [reported minimum central pressure (hPa) **929**]

5-10 10-20 20-30 30-40 40-50 50-60 60-70 70-80 80-90 >90%




Mean Sea Level Pressure in Tropical Cyclone Centre (hPa) **solid**=HRES; **dot**=Ens Mean




Date 20170905 12 UTC @ ECMF

Probability that **IRMA** will pass within 120 km radius during the next 240 h
tracks: **solid**=HRES; **dot**=Ens Mean [reported minimum central pressure (hPa) **929**]

5-10 10-20 20-30 30-40 40-50 50-60 60-70 70-80 80-90 >90%



Mean Sea Level Pressure in Tropical Cyclone Centre (hPa) **solid**=HRES; **dot**=Ens Mean



Question 1

What are some current advantages of using single-model ensembles?

- A. Estimates of uncertainty
- B. TC intensity model spread
- C. Alternative TC-track solutions
- D. All of the above
- E. A & C

Online Access to Ensemble Output and Training Resources

Access to Ensemble Output

NCEP GEFS and NAEFS: <http://mag.ncep.noaa.gov/>

National Weather Service
NCEP Central Operations

Home News Organization Search Search

Local forecast by "City, St" Go
Search NCEP Go

NCEP Quarterly Newsletter

Current Hazards
Watches/Warnings
Outlooks
National
Current Conditions
Observations
Satellite Images
Radar Imagery
Lakes & Rivers
Space Weather
Unified Surface
Analysis
Northern Hemisphere
Surface Analysis
Product Loops
Environmental Models
Product Info
Current Status
Model Analyses & Guidance
Forecasts
Current
6 to 10 Day
Aviation
Hurricane
Marine
Tropical Marine
Fire Weather
Forecast Maps
Climate
Climate Prediction
Climate Archives
Weather Safety
Storm Ready
NOAA
Central Library
Photo Library

Model Guidance

Back Home

Reset Selection(s)

Choose a Model Area or re-select a different Model Type

| Model Area | NAMER | SAMER | AFRICA | NPAC | EPAC | WNATL |
|------------|-------|--------|--------|------|--------|-------|
| ATLANTIC | POLAR | ATLPAC | EUS | WUS | ALASKA | |
| EUROPE | ASIA | SPAC | ARCTIC | | | |


| Model Type | GFS | NAM | SREF | VWV3 | HRW-NMM-EUS | HRW-ARW-EUS |
|-------------|-----------|-------|----------|-------------|-------------|-------------|
| GEFS-SPAG | NAM-HIRES | NAEFS | VWV3-ENP | HRW-NMM-WUS | HRW-ARW-WUS | |
| GEFS-MNSPRD | RAP | POLAR | VWV3-WNA | HRW-NMM-AK | HRW-ARW-AK | |

MAG v3.2.0

NOAA/ National Weather Service
National Centers for Environmental Prediction
5830 University Research Court
College Park, MD 20740
NCEP Internet Services Team
Page last modified: September 30 2013 18:48 PM UTC.

Disclaimer
Credits
Glossary


Privacy Policy
About Us
Career Opportunities



Access to ensemble
mean, spread, and
spaghetti plots


Canadian Ensembles

http://weather.gc.ca/ensemble/index_e.html




Spaghetti diagram of 500-mb 522 and 582 dm height contours

FHR 72 forecast of the probability that the 12 hour accumulation exceeds 2 mm
(The 12-h accumulation period immediately precedes the valid time)



Access to Ensemble Output

- ECMWF Ensembles:
<https://www.ecmwf.int/en/forecasts/charts>



COMET Courses

<http://www.meted.ucar.edu>

- Introduction to Ensemble Prediction:
http://www.meted.ucar.edu/nwp/pcu1/ensemble_webcast/
- Ensemble Forecasting Explained:
<http://www.meted.ucar.edu/nwp/pcu1/ensemble/>
- Ensemble Prediction System Matrix: Characteristics of Operational Ensemble Prediction Systems (EPS):
http://www.meted.ucar.edu/nwp/pcu2/ens_matrix/
- Wave Ensembles in the Marine Forecast Process:
<http://www.meted.ucar.edu/nwp/WaveEnsembles/>
- NWP Workshop on WRF and NAEFS:
http://www.meted.ucar.edu/s_africa_work/

Thank you

Questions?